


***Myzus persicae* neonicotinoid resistance management guidelines for Stone Fruits in Southern Europe, IRAC SPWG, 2014**

This is an update of the resistance alert and management recommendations issued in January 2013 by the IRAC Sucking Pest Working Group. It concerns specifically the appearance of neonicotinoid resistance in green peach aphid (*Myzus persicae*) in the peach orchards of southern France and north-eastern Spain and Northern Italy in 2010. The resistance is based on a target-site mutation which strongly affects neonicotinoid efficacy^{1,2}. The results of surveys from 2010 to 2012 confirmed the spread and presence of neonicotinoid-resistant aphids in many of the stone fruit orchards of southern France, Spain and Italy^{3,4}.

Map of the region showing areas where target site resistance to neonicotinoids was detected in *Myzus persicae* collected from stone fruit orchards from 2010 to 2012. No reports from new regions have been received to date in 2013.



IRAC have worked with local agricultural ministry officials, and entomological experts from Spain, France, Italy and the UK, to provide the following advice for the 2014 season in stone fruits, notably peaches:

Where no loss of performance to neonicotinoids has been experienced, it is recommended to use a maximum of one neonicotinoid application per crop cycle against *Myzus persicae* to minimise the further spread and intensification of the resistance and maintain effectiveness of the neonicotinoids. Depending on crop and country and local guidelines, this single spray may be pre-flowering or post-flowering, to fit with local label recommendations. (Note: Following restrictions to the neonicotinoids imidacloprid, thiamethoxam and clothianidin announced in 2013 by the European Commission, the recommended rotation programme has been modified accordingly to comply with these restrictions. See attached rotation scheme)

It is recommended that growers that have experienced a decline in activity to neonicotinoids in past seasons do not continue to use this group of insecticides as a preventative measure to halt the spread of resistance, and use insecticides with other modes of action, according to local registrations, such as products from groups 1A, 3A, 9B, 9C, 23⁵ as well as mineral oil to control *Myzus persicae*⁶. IRAC supports the use of any other IPM measures locally recommended, and may assist with the characterisation of resistance mechanisms in local *Myzus* populations⁷.

Acknowledgements: Many thanks to representatives of Rothamsted Research International, Università Cattolica del Sacro Cuore - Piacenza campus, Italy, University of Cartagena, Spain, Chamber of Agriculture in Catalunya and Aragon in Spain and the IRAC Spain Sucking Pest Working Group for inputs into these IRM recommendations.

- ¹ Bass et al. BMC Neuroscience 2011, 12:51 Mutation of a nicotinic acetylcholine receptor beta subunit is associated with resistance to neonicotinoid insecticides in the aphid *Myzus persicae*
- ² Puinean et al Pest Manag Sci (2012), Society of Chemical Industry, Development of a high-throughput real-time PCR assay for the detection of the R81T mutation in the nicotinic acetylcholine receptor of neonicotinoid-resistant *Myzus persicae*.
- ³ Slater et al, Pest Manag Sci 2012; 68: 634–638, Society of Chemical Industry, Identifying the presence of neonicotinoid resistant peach-potato aphid (*Myzus persicae*) in the peach-growing regions of southern France and northern Spain
- ⁴ Panini et al, Pest Manag Sci (2013), Society of Chemical Industry, Detecting the presence of target-site resistance to neonicotinoids and pyrethroids in Italian populations of *Myzus persicae*
- ⁵ See IRAC guidelines on *Myzus persicae* resistance management on the IRAC website
- ⁶ Consult local advisors for advice on which aphicides are affected by resistance in your locality.
- ⁷ Contact IRAC Sucking Pest Working Group or IRAC Spain on www.irac-online.org

Example of an IRM-based program in Stone Fruits in Southern Europe to limit spread of target site neonicotinoid resistance in *Myzus persicae*:

IRAC management recommendations for neonicotinoid resistant *Myzus persicae*:
Example 2014: Peaches, Nectarines in Southern Europe

